

## PROXIMATE COMPOSITION OF THREE COMMERCIAL FISHES COMMONLY CONSUMED IN AKWA IBOM STATE, NIGERIA

Daniel, Imaobong E.

Department of Chemistry, University of Uyo, Akwa Ibom State  
NIGERIA

### ABSTRACT

Proximate compositions of *Cynoglossus senegalensis*, *Polydactylus quadrifilis* and *Chrysichthys nigrodigitatus* obtained from Ayadehe in Akwa Ibom State, Nigeria were determined using standard methods. Results of the mean values for moisture, protein and fat were as follows; 66.19%, 14.2% and 10.62% respectively in *Chrysichthys nigrodigitatus*, 63.97%, 18.56% and 2.57% in *Cynoglossus senegalensis*, and 67.54%, 17.83% and 3.39% in *Polydactylus quadrifilis*. The ranges of ash carbohydrate, calorific value and crude fibre were as follows; 5.3% – 7.7%, 3.69% – 7.26%, 152.0 kcal/100g – 189.02 kcal/100g and 7.67% - 10.71% respectively. The data showed that these species of fishes are rich in crude protein, lipid, moisture and ash and meet the requirement for human nutritional needs.

**Keywords:** Calorific value, moisture, proximate composition.

### INTRODUCTION

Fish is one of the most important animal protein and other vital nutrients sources that are widely consumed by all races and classes of people (Abolude and Abdullahi, 2005; Andrew, 2001). Fish meat contains significantly low lipids and high water compared to that of beef or chicken and is favored over other white or red meats (Nestel, 2000). Lipids from fish are well known as a rich source of long-chain n-3 polyunsaturated fatty acids (LC n-3 PUFA) such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) which cannot be synthesized by humans and commonly obtained from the diet (Alasalvar *et al.*, 2002). Polyunsaturated fatty acids from fish have been reported to have preventive and/or curative effects for several diseases including arterial hypertension cancers and inflammatory diseases (Turkmen *et al.*, 2005). It may also aid in lowering the risk of Dementia, Alzheimer's diseases (Grant 1997), and prevent the cardiovascular diseases (Cahu *et al.*, 2004). Proximate composition is a good indicator of physiology which is needed for routine analysis of fisheries (Cui and Wootton, 1988). However, fish of various species do not provide the same nutrient profile to their consumers (Soriguer *et al.*, 1997; Takama *et al.*, 1999). These differences in the nutritional compositions of different species may be attributed to food composition, food and feeding habit, feeding rate, habitats, sex, age, size, genetic traits and season/migration (Dawson and Griman, 1980; Abdullahi, 2001; Ajah, 2009). Besides being used as food, fish is also increasingly demanded for use as feed. However, information concerning the chemical composition of freshwater fishes in general is valuable to nutritionists concerned with readily available sources of low-fat, high-protein foods such as most freshwater fishes (Sadiku and Oladimeji, 1991; Mozaffarian *et al.*, 2003; Foran *et al.*, 2005) and to the food scientist who is interested in developing them into high-protein foods, while ensuring the finest quality flavor, color, odor, texture, and safety obtainable with maximum nutritive value (Elagba *et al.* 2010). The aim of this study is to compare the nutritional values of three selected different fishes, in order to help consumers in choosing fish based on their nutrient values.

## MATERIALS AND METHODS

Fish species selected for this study were *Cynoglossus senegalensis*. (Tongue fish), *Polydactylus quadrifilis* (African giant threadfin), and *Chrysichthys nigrodigitatus* (silver catfish). These species were chosen for the study because of their abundance, high preference by people of Akwa Ibom State and market value. Eighteen (18) fresh fish samples were obtained directly from the fishermen on landing at Ayadehe in Akwa Ibom State. These comprised of six samples of *Chrysichthys nigrodigitalus* six samples of *Cynoglossus senegalensis* as well as six samples of *Polydactylus quadrifilis*. The collected samples were immediately packed into an ice chest to sustain freshness, and taken to the laboratory. Upon arrival to laboratory, the fishes were de-scaled where necessary, beheaded, gutted thoroughly washed with distilled water and filleted. Fillets of the same species were pooled together to form composite samples. These composite fillets were packed in labeled foil bowls, and kept frozen until needed for analysis.

### Proximate determination

Each fish sample in its fresh state was subjected to chemical analysis in triplicate according to AOAC 1990. Water content was determined by drying samples at  $105 \pm 2^{\circ}\text{C}$  until a constant weight was obtained. Dried samples were used for determination of crude fat, protein and ash contents. Crude fat was measured by solvent extraction method in a soxhlet system where petroleum ether was used as solvent. Crude protein content was calculated by using nitrogen content obtained by Kjeldahl method. A conversion factor of 6.25 was used for calculation of protein content. Crude fibre by acid base- digestion using 1.25%  $\text{H}_2\text{SO}_4$  (W/V) and 1.25% NaOH(W/V) solution. Carbohydrate content value were calculated based on difference calculation [Carbohydrate =  $100\% - (\% \text{moisture} + \% \text{ash} + \% \text{crude protein} + \% \text{fat})$ ], while the energy value was determined indirectly using Rubner's coefficients for aquatic organisms: The caloric values of the samples was obtained by multiplying the value of the crude protein, lipids and carbohydrates by 5.5, 9.5 and 4.1kcal respectively and taking the sum of the product (Winberg 1971).

### Data analysis

Date analysis was done using Analysis of variance (ANOVA) according to the statistical analysis system (Steel and Torie, 1987). Differences among sample means were tested for significance with Duncan's multiple range test (Duncan, 1955) at a level of 0.05

## RESULTS

Table 1: \*The proximate composition of *Cynoglossus senegalensis*, *Polydactylus quadrifilis* and *Chrysichthys nigrodigitatus* from Ayadehe in Itu LGA.

% composition	<i>C. nigrodigitatus</i>	<i>C. senegalensis</i>	<i>P. quadrifilis</i>
Moisture	66.19	63.97	67.54
Ash	5.3	7.7	5.94
Crude fibre	7.67	10.71	9.51
Crude protein	14.2	18.56	17.83
Fat	10.62	2.57	3.39
Carbohydrates	3.69	7.26	5.30
Calorific value(Kcal/100g)	189.02	155.94	152.0

\*Mean of three determinations.

The results of the nutritional values of the three species of fish selected for this study are presented in table 1. The moisture contents ranged from 63.97% in *Cynoglossus senegalensis* to 67.54% in *Polydactylus quadrifilis*. Ash content was lowest in *Chrysichthys nigrodigitatus* (5.3%), while *Cynoglossus senegalensis* had the highest % of ash (7.7%). Crude fibre ranged from 7.67% in *Chrysichthys nigrodigitatus* to 10.71% in *Cynoglossus senegalensis*, while crude protein ranged from 14.2% to 18.56% in *Chrysichthys nigrodigitatus* and *Cynoglossus senegalensis* respectively. For fats, *Chrysichthys nigrodigitatus* recorded the highest value of 10.62%, while the least value was recorded in *Cynoglossus senegalensis*. The carbohydrate content was between 3.69% (*Chrysichthys nigrodigitatus*) and 7.26% (*Cynoglossus senegalensis*). *Chrysichthys nigrodigitatus* had the highest energy value (189.02Kcal/100g), while the least was recorded for *Polydactylus quadrifilis*.

## DISCUSSION

The proximate composition of nutrients in all fish samples showed variation among the individual species. The moisture contents in all the three species were within the acceptable level (60-80 %), which could be due to the stable water levels in the location where the fish were collected (Adewumi *et al.*, 2014). The percentage of water is also a good indicator of its relative content of energy, protein and lipid (Olagunju., *et al.* 2012). Moisture content in all the species agreed with observation of Udo and Arazu (2012), Olagunju *et al.*, (2012), and Mazumder *et al.*, (2008) in several freshwater fish species. There were variations in the protein content of the three species examined in this study, with *Cynoglossus senegalensis* recording the highest value. The high protein content of *Cynoglossus senegalensis* may be attributed to the fact that they are carnivorous and known to feed on mollusks, shrimps, crabs and fish. Abdullahi (2001) reported that the protein content in fish vary with species due to certain factors such as the season of the year, effect of spawning and migration, food availability etc. However, despite the variations, the range of protein in fish in this study indicates that these species of fishes are good sources of protein to consumers. According to Ackman [1989], fish can be grouped into four categories according to their fat content: lean fish (< 2 %), low fat (2 to 4 %), medium fat (4 to 8%), and high fat (> 8%).

From the results obtained, the mean lipid contents in *Polydactylus quadrifilis* and *Cynoglossus senegalensis* indicates that they are low fat fish, while *Chrysichthys nigrodigitatus* can be classified as a high fat fish. This indicates that *Chrysichthys nigrodigitatus* are better sources of lipid in the body when consumed. Udo (2012) equally reported high values of fat for *Heterobranchus longifilis*, *Clarias gariepinus* and *Chrysichthys nigrodigitatus* of the Cross River, Nigeria. However, this result is at variance with 2.13% protein contents for *C. nigrodigitatus* obtained from Yenegoa as reported by Keremah and Amakiri (2013). The differences in these values could be due to the fact that these species were from different locations. Age variation, season and maturity in the same species may also contribute to the significant differences in the total lipid (Piggot and Tucker, 1990). Ash is a measure of the mineral content of food item. It is the inorganic residue that remains after the organic matter has been burnt off (Adewumi *et al.*, 2014). The range of ash in this study suggests that these species of fishes are good source of minerals such as calcium, potassium, zinc, iron and magnesium. Carbohydrates are sources of instant energy, which can be used in the body's development and growth (Olagunju *et al.*, 2012). Fish generally have very low levels of carbohydrates because glycogen does not contribute much to the reserves in the fish body tissue (USDA (2010; Jayasree *et al.*, 1994; Das and Sahu 2001). Carbohydrate content of fishes in this study was above known values of carbohydrates as reported by Vlieg (1988). The calorific value (kcal/100g) measured in *C. nigrodigitatus*, was

significantly higher than that determined in *Cynoglossus senegalensis* and *Polydactylus quadrifilis*. The high calorific value could be attributed to high fat content observed in *C. nigrodigitatus* in this study (Keremah and Amakiri 2013). This presumes that *C. nigrodigitatus* probably contain more calories than that available in the other two species.

## CONCLUSION

This study revealed the importance of *C. nigrodigitatus*, *C. senegalensis* and *p. quadrifilis* as good sources of protein and other nutrients. On the other hand, the information will be useful to the consumers in choosing fish based on their nutritional values rather than taste, appearance and other physical features.

## REFERENCES

- Abdullahi, S.A., (2001). Investigation on nutritional status of *Chrysiichthys nigrodigitatus*, *Barus filamentous* and *Aucheoglanis occidentalis* (Family: Bagridae). *Journal of Arid Zone Fisheries*, 1: 39-50
- Abolude, D S, & Abdullahi, S. A. (2005). Proximate and mineral contents in component parts of *Clarias gariepinus* and *Synodontis schall* from Zaria, Nigeria. *Nigerian Food Journal* 23:1- 8.
- Ackman RG. (1989). Nutritional composition of fats in seafoods. *Progress in Food Nutrition Science*. 13:161-241
- Adewumi, A.A, Adewole, H.A & Olaleye V.F (2014). Proximate and elemental composition of the fillets of some fish species in Osinmo Reservoir, Nigeria. *Agriculture and biology journal of North America*, 5(3): 109-117
- Ajah, P.O., (2009). *Fish breeding and hatchery management*. Jerry commercial productions, Calabar, p: 183.
- Alasalvar, C. Taylor, K.D., Zubcov, E., Shahidi, F.& Alexis, M. (2002). Differentiation of cultured and wild sea bass (*Dicentrarchus labrax*): total lipid content, fatty acid and trace mineral composition. *Food Chemistry*; 79:145 150.
- Andrew, A.E (2001): *Fish Processing Technology*. University of Ilorin press, Nigeria, pp 7-8.
- AOAC. 1990. In: Helrich, K (Ed.), Official Methods of Analysis of the Association of Official Analytical Chemists (15th edition). Section 969.33. Airlington: Association of Official Analytical Chemists.
- Cahu C, Salen E, & Lorgeril MD (2004). Farmed and wild fish in the prevention of cardiovascular diseases: Assessing possible differences in lipid nutritional values. *Nutrition Metabolism and Cardiovascular Diseases* 14: 34-41.
- Cui, Y. & Wooth, R.,J. (1988): Effect of ration, temperature and body size on the body composition, energy content and condition of Minnow (*Phoxinus phoxinus* L)
- Das, S., & Sahu, B.,K. (2001). Biochemical composition and calorific content of fishes and shellfishes from Rushikulya estuary, south Orissa coast of India. *Indian Journal of Fisheries* 48: 297-302.
- Dawson, A.S. & Griman, A.S. (1980). Quantitative seasonal changes in the protein, lipid and energy content of carcass, ovaries and liver of adult plaice (*Pleuronectes platena*). *Journal of Fish Biology*, 16: 943-945.
- Duncan D B (1955). Multiple Ranges and Multiple F-test. *Biometrics* 11:1-42
- Elagba, H. A. M., Rabie, A. and Mansour, H. M.(2010). Proximate composition, amino acid and mineral contents of five commercial Nile fishes in Sudan. *African Journal of Food Science* 4(10): 650-654
- Foran, J.A, Carpenter, D.O, Hamilton, M. C, Knuth, B. A, & Schwager, S. J. (2005). Risk-

- based consumption advice for farmed Atlantic and wild Pacific salmon contaminated with dioxins and dioxin-like compounds. *Environmental Health Perspectives.*, 33: 552-556.
- Grant, W.B. (1997) Dietary links to Alzheimer's disease. *Alzheimer's Disease Review* 2: 42-55.
- Jayasree V, Panilekar AH, Wahidulla S, Kamat SY (1994) Seasonal changes in biochemical composition of *Holothuria leucospilota* (Echinodermata). *Indian Journal of Marine Science* 23: 117-119.
- Keremah R. I. and Amakiri G.(2013). Proximate composition of nutrients in fresh adult catfishes: *Chrysichthys nigrodigitatus*, *Heterobranchus bidorsalis* and *Clarias gariepinus* in Yenagoa, Nigeria. *Greener Journal of Agricultural Sciences*, 3 (4): 291-294.
- Mazumder, M. S. A., Rahman, M. M., Ahmed, A. T. A., Begum, M. & Hossain, M. A. (2008). Proximate Composition of Some Small Indigenous Fish Species (SIS) in Bangladesh. *International Journal of Sustainable Crop Production*. 3(3):18-23
- Mozaffarian, M.,D, Rozenn, N.L, Lewis, H. K, Gregory, L.B, Russell, P.T, & Davis, S.S. (2003). Cardiac benefits of fish consumption may depend on type of fish meal consumed. *Circulation*. 107 : 1372-1382.
- Nestel, P.J. N. (2000). Fish oil and cardiovascular disease: lipids and arterial function. *American Journal of Clinical Nutrition*. 71:228-231
- Olagunju, A, Muhammad, A, Mada, S.B, Mohammed, A, Mohammed H. M, & Mahmoud, K. T.(2012) Nutrient Composition of Tilapia zilli, Hemisynodontis membranacea, Clupea harengus and Scomber Scombrus Locally Consumed in Africa. *World Journal of Life Sciences and Medical Research*; 2:16 – 19
- Piggot, G.M. & Tucker, B.W. (1990). Seafood: Effects of Technology on Nutrition. New York, USA: Marcel Dekker, Inc.
- Sadiku, S.O.E, & Oladimeji A.A. (1991). Relationship of proximate composition of Lates niloticus (L), synodontis schall. *Research Communication.*, 3: 29- 40.
- Soriguer, F., Serna, S., Valverde, E. (1997). Lipid, protein, and calorie content of different Atlantic and Mediterranean fish, shellfish, and mollusks commonly eaten in the south of Spain. *European Journal of Epidemiology*, 13: 451–463.
- Steel, R.G. & Torie, J. A (1987). Principles and Procedures of Statistics. A Biochemical Approach. 2nd Edn. Mc Graw- Hill International, Auckland, pp. 5-102.
- Takama, K., Suzuki, T., Yoshida, K., Arai, H. & Mitsui, T. (1999). Phosphatidylcholine levels and their fatty acid compositions in teleost tissues and squid muscle. *Comparative Biochemistry and Physiology Part B*, 124, 109–116.
- Turkmen, A., Aro, T., Nurmi, T., & Kallio, H (2005). Heavy metals in three commercially valuable fish species from Iskenderun Bay. Northern East Mediterranean Sea, Turkey. *Food Chemistry*. 91:167-172.
- Udo, P. J. (2012). Investigation of the Biochemical Composition of *Heterobranchus longifilis*, *Clarias gariepinus* and *Chrysichthys nigrodigitatus* of the Cross River, Nigeria. *Pakistan Journal of Nutrition* 11 (10): 865-868
- USDA (2010) US Department of Agriculture, Agricultural Research Service, National Nutrient Database for standard reference, Release 23. Nutrient laboratory.
- Vlieg, P., (1988). The proximate composition of New Zealand Marine fin fish and Shell fish. FAO corporate Document Repository, New Zealand Institute for crop and Food research. ISBN-477- 02520-X
- Winberg GG (1971). Symbols, units and conversion factors in studies of fresh water Productivity, pp. 134-178.